

DO GAMMA-RAY BURST SOURCES REPEAT?

Ramesh Narayan and Tsvi Piran*

Harvard-Smithsonian Center for Astrophysics,
60 Garden Street, Cambridge, MA 02138, U.S.A.

Summary

Following the discovery by Quashnock and Lamb (1993) of an apparent excess of γ -ray burst pairs with small angular separations, we reanalyze the angular distribution of the bursts in the BATSE catalogue. We find that in addition to an excess of close pairs, there is also a comparable excess of antipodal bursts, i.e pairs of bursts separated by about 180 degrees in the sky. Both excesses have only modest statistical significance. We reject the hypothesis put forward by Quashnock and Lamb that burst sources are repeaters, since it is obvious that this hypothesis does not predict an excess of antipodal coincidences. Lacking any physical model of bursts that can explain the antipodal pairs, we suggest that the two excesses seen in the data are either due to an unusual statistical fluctuation or caused by some unknown selection effect.

keywords: Gamma-rays: bursts

1. Introduction

In a recent paper Quashnock and Lamb (1993, denoted QL) analyze the distribution of angular separations of bursts in the publicly available BATSE catalogue of gamma-ray bursts. They find a significant excess of close pairs of bursts with angular separations smaller than $\sim 4^\circ$, compared to the number of such pairs expected for a random distribution of positions on the sky. On this basis they suggest that γ -ray bursters repeat. In their hypothesis, the close pairs actually arise from the same source, but they are assigned slightly different positions because of measurement errors, which are typically about 4° or larger. If QL are correct, then many extragalactic models may be ruled out; in particular, the neutron star merger model (Eichler et al, 1989; Narayan, Paczynski and Piran 1992) would become rather unlikely. Furthermore, as QL argue, their result implies a close relationship between classical γ -ray bursts and the three soft γ -ray repeaters. Since the latter are known to be located close to or in the Galaxy, this would give further support to the suggestion that the classical burst sources too are located in the Galaxy. In view of the importance of these conclusions we reanalyze in this letter the angular distribution of gamma ray bursts.

We repeat here the nearest neighbour analysis employed by QL and add to it another analysis based on the more standard angular autocorrelation function. We find that, while there does appear to be an excess of close pairs of bursts with angular separations less than 4° as claimed by QL, there is also an excess of antipodal pairs of bursts with angular

* Permanent Address: Racah Institute for Physics, The Hebrew University, Jerusalem 91904, Israel

separations larger than 176° . QL explain the excess of nearby bursts as signals from repeating sources, but we cannot think of any physical model to explain the excess of antipodal bursts. We thus conclude that both effects, if real, are most likely due to some unknown selection effect.

In sec. 2 of this letter we analyze the data. We examine the correlation function of the full sample of 260 bursts in the BATSE catalogue in sec. 2.a. and we present the nearest and farthest neighbour analysis for this sample in sec. 2.b. QL do not use the full sample, but analyze a subsample defined by the availability of counts in both the 64 ms and the 1024 ms channels on BATSE. Another subset could be defined by including only those bursts for which the formal positional accuracy is better than 4° . We discuss results from these subsamples in sec. 2.c. We conclude in sec. 3 with a discussion of the implications of the results.

2. Analysis

2.a The Two Point Angular Correlation Function

A data set composed of randomly positioned sources, some of which repeat, has a simple angular correlation function: a delta function peak at the origin due to the repeaters, and a slightly negative constant elsewhere. Errors in position measurements will spread out the delta function to a broadened peak at the origin with a width and shape corresponding to the probability distribution of the position errors, but nowhere else do we expect any other significant peak or dip. This simple structure of the two-point correlation function suggests that it ought to be a clean statistic to test the repeating source hypothesis.

The correlation function of all the 260 bursts in the BATSE catalogue is shown in Fig. 1(a). As expected from the QL analysis, there is a peak in the bin corresponding to $\theta < 4^\circ$. To assess the statistical significance of this peak we have carried out Monte Carlo simulations with 10000 random samples. From this we estimate that the peak has an amplitude of 1.75 standard deviations (1.75σ), and that the probability of obtaining a peak as strong as or stronger than the observed one is 0.0957 (9.57%). These results are given in Table 1 along with the results of various other calculations discussed later in the paper. Note that the probability distribution of the values of the correlation function is not Gaussian. Therefore, here and elsewhere in the text, all significance levels that we quote are obtained from Monte Carlo simulations. Note also that a small probability in Table 1 means that it is unlikely for the particular event to have happened by chance and therefore implies high significance.

While the existence of a peak at $\theta < 4^\circ$ agrees with the QL result, there is another unexpected peak at the antipode, corresponding to the bin with $\theta > 176^\circ$. The second peak is slightly wider than the peak at the origin and has an amplitude of 1.86σ , which is marginally more significant (probability = 0.0724) than the direct peak. In fact, while the statistical significance of each of the single peaks may be considered marginal, the statistical significance of having *both* peaks is much higher. Additionally, there are weak negative excursions in the correlation function at around $\theta \sim 10 - 20^\circ$ and $\theta \sim 160 - 170^\circ$ which appear to be moderately significant, but we do not discuss these in detail.

2.b Nearest Neighbour Analysis.

Quashnock and Lamb did not employ the correlation function but instead used a “nearest neighbour statistic.” According to this statistic they measure for each burst in the catalogue the angular distance to its nearest neighbour and compute the cumulative distribution of this quantity for the observed sample. They then compare this cumulative with the distribution expected for a random sample. In Fig. 1(b) we present this comparison for the full sample of 260 bursts. We show the number of bursts with nearest neighbours closer than an angle θ as a function of $1 - \cos \theta$. As expected from QL’s analysis and from the correlation function approach discussed in sec. 2.a., there is evidence for an excess of nearby bursts. Motivated by our discovery of the antipodal peak in the correlation function, we also calculate for each burst the distance to its *farthest* neighbour, that is the burst that is nearest to the antipodal point. We plot this cumulative distribution in Fig 1(b), where now the x-axis is $1 + \cos \theta$ and the vertical axis represents the number of bursts with farthest neighbour more distant than angle θ . As expected from the correlation function, we find that there is again an excess of antipodal bursts compared to the distribution expected for random bursts. Visually at least, we would say that the evidence for an excess is about equally strong for nearest and farthest neighbours.

What is the significance of these deviations? Since Fig. 1(b) is a cumulative distribution, one might consider using the Kolmogorov-Smirnov (KS) test to estimate the significance. In this method one measures the KS distance, which is the maximum distance between the observed cumulative curve and the theoretically expected curve. One then calculates the probability of obtaining a distance at least as large as the observed one by the standard KS method (e.g. Press et al, 1992). However, the KS method assumes that all the data points are independent, but this is not the case in the present distribution because the set of nearest neighbour distances can have strong correlations. To see this, consider the case when two bursts happen to lie very close to each other on the sky. Both bursts will yield the same small nearest neighbour distance, and therefore this distance will be counted twice. This demonstrates that a direct application of the KS method is invalid.

To measure correctly the statistical significance of the deviations we generated 10000 random data sets and calculated the distribution of the KS distance. We find that the probability of obtaining KS distances greater than those observed in the 260 burst BATSE sample are: 0.012 for the nearest neighbours and 0.11 for the farthest neighbours. (Note that the standard KS test gives probabilities of 0.0016 and 0.019, showing that the KS test, by neglecting the effect of correlations, leads to unduly optimistic estimates of the significance.) The results are marginally significant at best. If we estimate the probability for both nearest and farthest neighbours to have such large deviations simultaneously, we find that it is 0.0019, which is somewhat more significant. Therefore, once again we reach the same conclusion as we did from the correlation analysis in sec. 2.a, viz. while we find an excess of nearest and farthest neighbours in the data, these excesses are only marginally significant. However, the probability of obtaining both excesses simultaneously is very small and therefore the evidence for such a signal is more significant.

2.c Subsamples of the BATSE Catalogue

The basic variables in our analysis are the positions of the bursts. As stated in the instructions with the BATSE catalogue, burst positions have variable errors depending on the strengths of the bursts, the position and orientation of the Compton GRO satellite, and other parameters. The BATSE catalogue gives an estimate of the formal positional error $\Delta\theta_P$ for each burst due to Poisson fluctuations in the observed γ -ray counts, to which an additional systematic error of 4° should be added in quadrature to yield the total positional error, $\Delta\theta_{tot} = [\Delta\theta_P^2 + (4^\circ)^2]^{1/2}$. In some cases the estimated $\Delta\theta_P$ is quite large (up to 20°) and it is reasonable to exclude such bursts from the samples. Since we are looking for correlations on angular separations of 4° or less, it is clearly meaningless to argue that a burst with a positional error of say, 10° , is within 4° from another burst. We have therefore repeated our analysis with a sub sample of bursts for which the quoted formal $\Delta\theta_P$ values are 4° or less. This reduces the sample from 260 bursts to 131 bursts. When we analyze this sample, the correlation analysis gives a modest peak of 0.84σ for $\theta < 4^\circ$, and an impressive peak of 2.69σ for antipodal neighbours between 176° and 180° . The corresponding probabilities for chance occurrence are: 0.42 for the direct peak and 0.010 for the antipodal peak (see Table 1). Similarly, the QL statistic based on the KS distance gives excesses of 0.83σ and 1.61σ for nearest and farthest neighbours, corresponding to chance probabilities of 0.47 and 0.027. The fact that the statistical significance of the effect does not increase, and in fact *decreases*, when we throw away the bursts with large positional errors supports our suspicion that the observed anomaly is not due to a real physical effect.

For completeness, we have also repeated the analysis with the particular sub-sample used by QL. They divide the data into subgroups according to the γ -ray counts measured in the 64 ms and 1024 ms channels. Their total sample, identified as Type I+II in their paper, consists of 201 bursts; we refer to this as the QL sample. Among all the subsamples that QL analyzed, they found the strongest signal in this particular combined sample. Using 10000 Monte Carlo simulations with synthetic data, we find that there is a probability of 0.0015 of obtaining by chance a nearest neighbour deviation comparable to the signal observed in the QL sample. We also find a modest excess of farthest neighbours in this sample, with a Monte Carlo probability of 0.36.

Our final subset is a truncated QL sample of 108 bursts, where we take the QL sample and eliminate all bursts with $\Delta\theta_P > 4^\circ$. The hypothesis of repeating bursts put forward by QL suggests that by eliminating bursts with highly uncertain positions the statistical significance of the effect should increase. We find this not to be the case. The angular correlation function for this data set is shown in Fig. 2(a). We see both the forward and antipodal peaks, but the forward peak represents only a $.74\sigma$ deviation while the antipodal peak is at 2.60σ (see Table 1 for the corresponding probabilities). Similarly, we show in Fig. 2(b) the nearest/farthest statistic. Here we find that the KS distance for the distribution of nearest bursts is only 0.68σ (random chance probability of 0.66), while for the farthest bursts the KS distance is 1.24σ corresponding to a random chance probability of 0.10.

3. Summary and Discussion

Our primary conclusions are the following:

(i) The BATSE data does contain some evidence for an excess of pairs of bursts with angular separations smaller than 4° . However, there is equally good evidence for an excess of nearly antipodal pairs, with separations between 176° and 180° . An antipodal peak is, of course, not expected from a random population of repeaters.

(ii) The statistical significance of either excess is marginal. We find that the only statistic that provides a reasonably strong signal consistently is when we combine the excesses in the forward and antipodal directions and compute the probability of obtaining both simultaneously. There are obvious dangers in carefully selecting two hypotheses in this fashion and combining them, therefore we are ambivalent about accepting the combined excess as a real property of the BATSE sample of bursts.

(iii) When we eliminate bursts with Poisson positional errors $\Delta\theta_P > 4^\circ$ (which corresponds to total error $\Delta\theta_{tot} > 5.7^\circ$), far from becoming stronger, the evidence for the excesses actually becomes weaker. In fact the decrease in the signal is quite drastic in the case of nearest neighbour pairs, while it is more modest for the antipodal pairs. In other words, the evidence for an excess of close pairs is very volatile, depending on the particular sample chosen, while the excess of antipodal pairs shows a little more stability at least within the tests we have done (Table 1). Note that bursts with large $\Delta\theta_P$ are generally weaker and one could argue that by eliminating them we eliminate the repeating weak bursts (according to the QL model). However, the large positional error of these weak bursts means that these bursts should not have shown any evidence for the repeater model in the first place.

We have been unable to come up with any physical model that can explain an excess of bursts within 4° of the antipode. Such an excess may occur if the bursts are located along a narrow line in space or in a very thin disk. However, such distributions are clearly ruled out by the observed overall isotropy of the positions of the bursts (Meegan et al, 1992).

We are forced to conclude that the excess of close pairs of bursts discovered by QL, and the antipodal excess that we discuss here, are either due to a statistical fluctuation or caused by some selection effect¶. We do not have any specific idea as to the nature of the selection effect.

We thank Dalia Goldwirth for useful discussions, and Don Lamb and Jean Quashnock for help in understanding their burst samples and for comments. This work was supported by NASA grant NAG 5-1904.

References

- Eichler, D., Livio, M., Piran, T. & Schramm, D.N. 1989, *Nature*, **340**, 126.
 Meegan, C.A., *et. al.* 1992, *Nature*, **355** 143.
 Narayan, R., Paczyński, B. & Piran, T. 1992, *ApJL*, **395**, L83.

¶ The possibility of a selection effect is supported by the fact that the two bursts identified by trigger numbers 803 and 974 in the BATSE catalogue have identical positions to all digits given. However, one of the bursts has a positional accuracy of only 16° , making the agreement of the two positions highly unlikely.

Press, W.H., Teukolsky, S.A., Vetterling, W.T. & Flannery, B.P. 1992, *Numerical Recipies*, Cambridge University Press, Cambridge.

Quashnock, J. Q & Lamb, D. 1993, MNRAS, submitted.

Table I- Correlations functions and significance levels

Sample	Number	$c(\leq 4^\circ)$	Prob	$c(\geq 176^\circ)$	Prob
Full Sample	260	$0.268 = 1.75\sigma$	0.096	$0.292 = 1.86\sigma$	0.072
QL Sample	201	$0.389 = 1.95\sigma$	0.052	$0.348 = 1.74\sigma$	0.082
Full, $\Delta\theta_P \leq 4^\circ$	131	$0.254 = 0.84\sigma$	0.423	$0.832 = 2.69\sigma$	0.010
QL, $\Delta\theta_P \leq 4^\circ$	108	$0.279 = 0.74\sigma$	0.575	$0.989 = 2.60\sigma$	0.016

Figure Captions

Fig. 1: (a) The angular autocorrelation function of the full sample of 260 bursts in the BATSE catalogue, represented in 4° bins. The error bars correspond to one standard deviation as estimated through Monte Carlo simulations. Note that the antipodal peak near $\theta \sim 180^\circ$ is somewhat higher and wider than the direct peak near $\theta \sim 0^\circ$. There are marginally significant negative dips adjacent to both peaks and it is plausible that the excess counts in the peaks are supplied by the lack of bursts in the dips. (b) The cumulative number of nearest neighbours closer than θ as a function of $1 - \cos \theta$ and farthest neighbours more distant than θ as a function of $1 + \cos \theta$, both shown as thick lines. The theoretically expected curve for a random sample is shown by the smooth thin line. Note that the excess extends to larger angles in the case of farthest neighbours, whereas the KS statistic (i.e. the largest distance between the observed and theoretical curves) is slightly higher in the case of nearest neighbours.

Fig. 2: (a) The angular autocorrelation function of the 108 bursts out of the QL sample which have Poisson position errors $\Delta\theta_P \leq 4^\circ$. Note that the antipodal peak is much stronger than the direct peak, the latter being practically insignificant. (b) The cumulative number of nearest neighbours as a function of $1 - \cos \theta$ and farthest neighbours as a function of $1 + \cos \theta$, shown as thick lines. The theoretically expected curve is shown by the thin line. The nearest neighbour curve is very close to the expected curve whereas the farthest neighbour curve shows an apparently significant deviation.